

Sandoz® Dimenhydrinate 100mg Adults 10 Dimenhydrinate Suppositories [1]



Recommended use or purpose:

Prevention and treatment of nausea, vomiting, dizziness and motion sickness.

Ingredients:

Medicinal ingredients: dimenhydrinate 100mg Non-medicinal ingredient: hard fat.

Dosage:

Adults: ½ or 1 suppository, 3 or 4 times a day as needed. Children 12 years and over: ½ suppository 2 or 3 times a day as needed. For motion sickness, dosage should be taken at least 30 min and preferably 1 or 2 hours before travelling.

Warning:

Do not exceed recommended dosage. May cause drowsiness, avoid driving a motor vehicle or engaging in activities requiring alertness. Avoid alcoholic beverages. Do not take if you have glaucoma, chronic lung disease, difficulty urinating due to an enlargement of the prostate gland, you are pregnant or breastfeeding, unless directed by a physician. Do not take with other antihistamines, tranquilizers or any other sedating drugs without first consulting a

physician. May cause excitability, especially in children. Prolonged use on advice of a physician only.

*For additional information, or to learn more about the proper use of this product, be sure to read the product leaflet and/or product monograph before use.

Product monograph ^[4]

® Registered trademark owned or used under license by Sandoz Canada Inc.

Accordion Type:

Collapsible

Source URL: <https://www.sandoz.ca/en/sandozr-dimenhydrinate-100mg-adults-10-dimenhydrinate-suppositories>

Links

[1] <https://www.sandoz.ca/en/sandozr-dimenhydrinate-100mg-adults-10-dimenhydrinate-suppositories>

[2] <https://pediavit.ca/en/>

[3] <https://www.sandoz.ca/sites/www.sandoz.ca/files/Omep%20Feuille%20EN.pdf>

[4]

https://www.sandoz.ca/sites/www.sandoz.ca/files/SUPPOSITORIES%20DIMENHYDRINATE_Product%20Monograph.pdf