

## **Sandoz® Dimenhydrinate 50mg Children 10** **Dimenhydrinate Suppositories** [1]



### **Recommended use or purpose:**

Prevention and treatment of nausea, vomiting, dizziness and motion sickness.

### **Ingredients:**

Medicinal ingredients: dimenhydrinate 50mg Non-medicinal ingredient: hard fat.

### **Dosage:**

Children 8 years and over: ½ or 1 suppository, 2 or 3 times a day as needed. For motion sickness, dosage should be taken at least 30 min and preferably 1 or 2 hours before travelling.

### **Warning:**

Do not exceed recommended dosage. May cause drowsiness, avoid driving a motor vehicle or engaging in activities requiring alertness. Avoid alcoholic beverages. Do not take if you have glaucoma or a chronic lung disease unless directed by a physician. Do not take with other antihistamines, tranquilizers or any other sedating drugs without first consulting a physician. May cause excitability, especially in children. Prolonged use on advice of a physician only.

\*For additional information, or to learn more about the proper use of this product, be sure to read the product leaflet and/or product monograph before use.

Product monograph <sup>[4]</sup>

® Registered trademark owned or used under license by Sandoz Canada Inc.

**Accordion Type:**

Collapsible

---

**Source URL:** <https://www.sandoz.ca/en/sandozr-dimenhydrinate-50mg-children-10-dimenhydrinate-suppositories>

**Links**

[1] <https://www.sandoz.ca/en/sandozr-dimenhydrinate-50mg-children-10-dimenhydrinate-suppositories>

[2] <https://pediavit.ca/en/>

[3] <https://www.sandoz.ca/sites/www.sandoz.ca/files/Omep%20Feuille%20EN.pdf>

[4] [https://www.sandoz.ca/sites/www.sandoz.ca/files/SUPPOSITORIES%20DIMENHYDRINATE\\_Product%20Monograph.pdf](https://www.sandoz.ca/sites/www.sandoz.ca/files/SUPPOSITORIES%20DIMENHYDRINATE_Product%20Monograph.pdf)