

PRODUCT MONOGRAPH

Pr CEFAZOLIN FOR INJECTION USP

cefazolin (as cefazolin sodium)

500 mg, 1 g & 10 g per vials

Antibiotic

STERILE

Sandoz Canada Inc.
110 rue de Lauzon
Boucherville, QC
J4B 1E6

Date of Revision:
October 04, 2018

Submission Control No: 219834

PRODUCT MONOGRAPH

Pr Cefazolin for Injection USP

THERAPEUTIC CLASSIFICATION Antibiotic

ACTION AND CLINICAL PHARMACOLOGY

Cefazolin sodium is a cephalosporin antibiotic for parenteral administration. Cefazolin exerts its bactericidal effect by inhibiting bacterial cell wall synthesis. Cefazolin is about 85% bound to serum protein. The peak level in serum is approximately 32-42 mg/mL after an intramuscular (IM) injection of 500 mg. Over 80% of injected cefazolin is excreted in the urine during the first 24 hours after IM injection; most is excreted during the first 4-6 hours.

INDICATIONS AND CLINICAL USE

Cefazolin for Injection USP (sterile cefazolin sodium) is indicated in the treatment of the following infections when caused by susceptible strains of the listed organisms:

RESPIRATORY TRACT INFECTIONS caused by *Streptococcus pneumoniae*, *Klebsiella pneumoniae*, *Hemophilus influenzae*, *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant) and group A *beta-hemolytic streptococci*.

URINARY TRACT INFECTIONS caused by *Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumoniae* and some strains of enterobacter, and enterococci. See NOTE below.

SKIN AND SOFT TISSUE INFECTIONS caused by *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant), group A *beta-hemolytic streptococci* and other strains of streptococci.

BONE AND JOINT INFECTIONS caused by *Staphylococcus aureus*.

SEPTICEMIA caused by *Streptococcus pneumoniae*, *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant), *Proteus mirabilis*, *Escherichia coli* and *Klebsiella pneumoniae*. See NOTE below.

ENDOCARDITIS caused by *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant) and group A *beta-hemolytic streptococci*.

Determine susceptibility of the causative organism to cefazolin sodium by performing appropriate culture and susceptibility studies should be performed. (See MICROBIOLOGY for disc susceptibility tests and dilution techniques).

NOTE: Most strains of *Enterococci*, indole positive *Proteus* (*P. vulgaris*), *Enterobacter cloacæ*, *Morganella morganii*, *Providencia rettgeri* and methicillin-resistant staphylococci are resistant. *Serratia*, *Pseudomonas*, and *Acinetobacter calcoaceticus* (formerly *Mima* and *Herellea* species) are almost uniformly resistant to cefazolin. (See MICROBIOLOGY).

Perioperative Prophylaxis: In patients undergoing potentially contaminated surgical procedures, and in patients in whom infection would pose a serious risk (e.g. during open-heart surgery and prosthetic arthroplasty), the preoperative, intraoperative and postoperative administration of Cefazolin for Injection USP may reduce the incidence of certain postoperative infections.

Identification of the causative organisms should be made by culture should signs of infection occur so that appropriate therapy may be instituted.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of CEFAZOLIN FOR INJECTION USP and other antibacterial drugs, CEFAZOLIN FOR INJECTION USP should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

Sterile cefazolin sodium is contraindicated in patients with known allergy to the cephalosporin group of antibiotics.

WARNINGS

Sterile cefazolin sodium should be used with caution in penicillin-allergic patients. There is clinical evidence of partial cross-allergenicity of the penicillins and the cephalosporins. There are instances of patients who have had reactions to both penicillins and cephalosporins (including fatal anaphylaxis after parenteral use). Clinical and laboratory evidence of partial cross-allergenicity of the two drug classes exists.

Sterile cefazolin sodium should be administered cautiously and then only when absolutely necessary to any patient who has demonstrated allergy, particularly to drugs. Immediate emergency treatment with epinephrine is indicated for serious anaphylactoid reactions. As indicated, oxygen, intravenous steroids, and airway management including intubation, should also be employed.

There have been reports of pseudo membranous colitis with the use of cephalosporins. It is therefore important to consider its diagnosis in patients who develop diarrhea in association with antibiotic use.

Susceptibility/Resistance

Development of Drug Resistant Bacteria

Prescribing CEFAZOLIN FOR INJECTION USP in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

PRECAUTIONS

The overgrowth of non-susceptible organisms may result from the prolonged use of sterile cefazolin sodium. It is essential that the patient be carefully observed. In patients with a history of lower gastrointestinal disease, particularly colitis, cefazolin sodium should be prescribed with caution.

Clinitest[®] tablets solution, but not enzyme-based tests such as Clinistix[®] and Tes-Tape[®] may falsely indicate glucose in the urine of patients on cefazolin.

Positive direct and indirect Coombs' tests have been reported during treatment with cefazolin. These may also occur in neonates whose mothers received cephalosporins before delivery. The clinical significance of this effect has not been established.

Use in Renal Impairment

Caution should be exercised in treating patients with pre-existing renal damage although cefazolin has not shown evidence of nephrotoxicity.

Patients with low urinary output due to impaired renal function should be administered reduced daily dosages of cefazolin. (See Dosage in Patients with Reduced Renal Function). Blood levels of cefazolin in dialysis patients remain fairly high and should be monitored.

Probenecid may decrease renal tubular secretion of cefazolin when used concurrently with sterile cefazolin sodium, resulting in increased and prolonged cefazolin blood levels.

In beta-hemolytic streptococcal infections, treatment should be continued for at least 10 days, to minimize possible complications associated with the disease.

Use in Pregnancy and Lactation

The safety of the use of cefazolin sodium during pregnancy has not been established.

Lactation

Very low concentrations of cefazolin are found in the milk of nursing mothers. Cefazolin sodium should be administered with caution to a nursing woman.

Children

The safety of the use of cefazolin sodium in prematures and infants under one month of age has not been established.

Drug Interactions

The renal tubular secretion of cefazolin may be decreased when probenecid is used concurrently, resulting in increased and prolonged cefazolin blood levels.

ADVERSE REACTIONS

The following reactions have been reported:

Gastrointestinal: Diarrhea, oral candidiasis (oral thrush), vomiting, nausea, stomach cramps and anorexia. During antibiotic treatment symptoms of pseudo membranous colitis can appear. There have been rare reports of nausea and vomiting.

Allergic: Allergic reactions occur infrequently and include: anaphylaxis, eosinophilia, itching, drug fever, skin rash.

Hematologic: Neutropenia, anemia, leukopenia, thrombocytopenia, positive direct and indirect antiglobulin (Coombs') tests.

Hepatic and Renal: Without clinical evidence of renal or hepatic impairment transient increases in AST (SGOT), ALT (SGPT), BUN and alkaline phosphatase levels have been observed.

Transient hepatitis and cholestatic jaundice have been reported rarely, as with some penicillins and some other cephalosporins.

Local Reactions: Phlebitis at the site of injection has occurred rarely. Infrequently there is pain at the site of injection following intramuscular injection. Some induration has been reported.

Other Reactions: Vulvar pruritus, genital moniliasis, vaginitis and anal pruritus.

SYMPTOMS AND TREATMENT OF OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

There is a lack of experience with acute sterile cefazolin sodium overdose. Supportive therapy should be instituted according to symptoms in cases of suspected overdose.

DOSAGE AND ADMINISTRATION

Dosage

After reconstitution Cefazolin for Injection USP may be administered either intramuscularly or intravenously. In both cases total daily dosages are the same.

Adults

Adult Dosage Guide

Type of Infection	Dose	Frequency
Mild infections caused by susceptible Gram-positive cocci	250 mg to 500 mg	Every 8 hours
Acute, uncomplicated urinary tract infections*	1 g	Every 12 hours
Moderate to severe infections	500 mg to 1 g	Every 6 to 8 hours

*This dosage recommendation applies to intramuscular use. The efficacy of cefazolin sodium when administered intravenously at 12 hour intervals has not been established.

Cefazolin for Injection USP has been administered in dosages of 6 g per day in serious infections such as endocarditis.

Treatment should be continued for at least 10 days in beta-hemolytic streptococcal infections to minimize possible complications associated with the disease.

Dosage in Patients with Reduced Renal Function

After an initial loading dose appropriate to the severity of the infection, the following reduced dosage schedule is recommended:

Dosage Guide for Patients with Renal Impairment

Creatinine Clearance (mL/s)	Serum Creatinine (mMol/L)	Dosage
≤0.91	≥ 140	250 mg to 1 g every 6-12 hours
0.58-0.90	141-273	250 mg to 1 g every 8-12 hours

0.18-0.57	274-406	125 mg to 500 mg every 12 hours
≤0.17	≥407	125 mg to 500 mg every 18 hours

Perioperative Prophylactic Use

The recommended dosage regimen to prevent postoperative infection in contaminated or potentially contaminated surgery is:

- One gram intravenously or intramuscularly administered ½ hour to 1 hour prior to the start of surgery so that at the time of the initial surgical incision adequate antibiotic levels are present in the serum and tissues.
- For lengthy operative procedures (e.g. 2 hours or more) 0.5-1.0 g administered intravenously or intramuscularly during surgery. (Administration should be modified according to the duration of the operative procedure and the time of greatest exposure to infective organisms.)
- Postoperatively, 0.5-1.0 gram intravenously or intramuscularly every 6 to 8 hours for 24 hours postoperatively. The prophylactic administration of cefazolin sodium may be continued for 3 to 5 days following the completion of surgery in which the occurrence of infection may be particularly devastating (e.g. open-heart surgery and prosthetic arthroplasty).

Children

A total daily dosage of 25 to 50 mg per kg (approximately 10 to 20 mg per pound) of body weight, divided into three or four equal doses, is effective for most mild to moderately severe infections in children.

For severe infections total daily dosage may be increased to 100 mg per kg (45 mg per pound) of body weight. The use of cefazolin in prematures and in infants under one month of age is not recommended since the safety for use in these patients has not been established.

Paediatric Dosage Guide – 25 mg/kg/day

Weight		25 mg/kg/day Divided Into 3 Doses		25 mg/kg/day Divided Into 4 Doses	
lb	kg	Approximate Single Dose mg/q8h	Volume Needed of 125 mg/mL* Solution	Approximate Single Dose mg/q6h	Volume Needed of 125 mg/mL* Solution
10	4.5	40 mg	0.35 mL	30 mg	0.25 mL
20	9.0	75 mg	0.60 mL	55 mg	0.45 mL
30	13.6	115 mg	0.90 mL	85 mg	0.70 mL

40	18.1	150 mg	1.20 mL	115 mg	0.90 mL
50	22.7	190 mg	1.50 mL	140 mg	1.10 mL

* 125 mg/mL concentration may be obtained by reconstituting the 500 mg vial with 3.8 mL of diluent.

Paediatric Dosage Guide – 50 mg/kg/day

Weight		50 mg/kg/day		50 mg/kg/day	
		Divided Into 3 Doses		Divided Into 4 Doses	
lb	kg	Approximate Single Dose mg/q8h	Volume Needed of 225 mg/mL* Solution	Approximate Single Dose mg/q6h	Volume Needed of 225 mg/mL* Solution
10	4.5	75 mg	0.35 mL	55 mg	0.25 mL
20	9.0	150 mg	0.70 mL	110 mg	0.50 mL
30	13.6	225 mg	1.00 mL	170 mg	0.75 mL
40	18.1	300 mg	1.35 mL	225 mg	1.00 mL
50	22.7	375 mg	1.70 mL	285 mg	1.25 mL

*225 mg/mL concentration may be obtained by reconstituting the 500 mg vial with 2 mL of diluent.

Treatment with 60 percent of the normal daily dose may be administered in divided doses every 12 hours to children with mild to moderate renal impairment (Ccr 0.67-1.17 mL/s). Children with moderate to severe renal impairment (Ccr 0.33-0.87 mL/s) should be given 25 percent of the normal daily dose in equally divided doses every 12 hours, and children with severe renal impairment (Ccr 0.08-0.33 mL/s) should receive 10 percent of the normal daily dose every 24 hours.

All dosage recommendations apply after an initial loading dose.

ADMINISTRATION

NOTE: See PHARMACEUTICAL INFORMATION for reconstitution and dilution directions.

Intramuscular Administration

Inject the reconstituted solution into a large muscle mass. Pain on injection of Cefazolin for Injection USP occurs infrequently.

Intravenous Administration

Direct (bolus) injection: Inject the appropriately diluted reconstituted solution slowly over 3 to 5 minutes directly into a vein or through tubing for patients receiving parenteral fluids. (See list

of solutions for intravenous infusion in PHARMACEUTICAL INFORMATION).

Intermittent or Continuous Infusion: The reconstituted solution can be administered along with primary intravenous fluid management programs in a volume control set or in a separate secondary IV bottle. (See list of solutions for intravenous infusion in PHARMACEUTICAL INFORMATION).

PHARMACEUTICAL INFORMATION


Drug Substance

Proper Name: Cefazolin sodium

Chemical Name: 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 3-[[[(5-methyl-1,3,4-thiadiazol-2-yl)thio]methyl]-8-oxo-7-[(1*H*-tetrazol-1-yl)acetyl] amino]-, monosodium salt (6*R*-*trans*).

Monosodium (6*R*, 7*R*)-3{[(5-methyl-1,3,4-thiadiazol-2-yl)thio]methyl}-8-oxo-7-[2-(1*H*-tetrazol-1-yl)acetamido]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate.

Structural Formula:



Molecular Formula: $C_{14}H_{13}N_8NaO_4S_3$

Molecular Mass: 476.50 g/mol

Description: Cefazolin sodium is a white to off-white, practically odourless solid. The drug is freely soluble in water and very slightly soluble in alcohol. It is practically insoluble in chloroform and ether. The pH is between 4.5 and 6.0, in an aqueous solution containing 100 mg of cefazolin per millilitre.

Composition

Each vial contains sterile cefazolin sodium. Each gram of cefazolin sodium contains 48 mg of sodium. Cefazolin for Injection USP does not contain any preservatives.

Stability and Storage Recommendations

Cefazolin for Injection USP (sterile cefazolin sodium) should be stored between 15 and 25°C, protected from light.

Reconstituted Solutions

Parenteral drug products should be SHAKEN TO DISSOLVE ALL POWDER when reconstituted, and inspected visually for particulate matter prior to administration. The drug solutions should be discarded if particulate matter is evident in reconstituted fluids.

Reconstituted solutions may range in colour from pale yellow to yellow without a change in potency.

Reconstituted Cefazolin for Injection USP may be stored for 24 hours between 15 and 25°C, or for 72 hours under refrigeration between 2 and 8°C, protected from light.

Cefazolin for Injection USP solution reconstituted with bacteriostatic diluent and used for intramuscular administration as multiple-dose containers should be used within 7 days when stored under refrigeration.

The Pharmacy Bulk Vial is intended for multiple dispensing for intravenous use only, employing a single puncture. Following reconstitution, the solution should be dispensed and diluted for use within eight hours. Any unused reconstituted solution should be discarded after eight hours.

(1) For Intramuscular Injection:

Single Dose Vials

Reconstitute according to the table which follows. SHAKE TO DISSOLVE ALL POWDER.

Single Dose Vial Reconstitution Table

Strength	Diluent	Volume to be Added to Vial (mL)	Approximate Available Volume (mL)	Nominal Concentration (mg/mL)
500	Sodium Chloride Injection	2.0	2.2	225
	OR Sterile Water for Injection	3.8	4.0	125
1 000	Sterile Water for Injection	2.5	3.0	334

(2) For Direct Intravenous (bolus) Injection:

Single Dose Vial

Reconstitute as directed above. SHAKE TO DISSOLVE ALL POWDER. A minimum of 10 mL of Sterile Water for Injection should be used to dilute the reconstituted solution.

Pharmacy Bulk Vial

Pharmacy Bulk Vials should be used for intravenous use only. Add, according to the table below, Sterile Water for injection, or Sodium Chloride injection. SHAKE TO DISSOLVE ALL POWDER.

Pharmacy Bulk Vial Reconstitution Table

Strength	Amount of Diluent	Approximate Available Volume	Approximate Concentration
10 grams	45 mL	50 mL	200 mg/mL
	96 mL	100 mL	100 mg/mL

*Shake to dissolve and let stand until clear.

The vial is intended for single puncture and multiple dispensing, and the vial contents should be used within 8 hours.

(3) For intermittent or continuous intravenous infusion, reconstituted Cefazolin for Injection USP may be further diluted as follows:

Single Dose Vials

Reconstitute according to the Single Dose Vial Reconstitution Table above. SHAKE TO DISSOLVE ALL POWDER. Further dilute the reconstituted Cefazolin for Injection USP to 50 to 100 mL in one of the following solutions:

- Sodium Chloride Injection 0.9%
- Dextrose Injection 5% or 10%
- Dextrose 5% in Lactated Ringer's Injection
- Dextrose 5% and Sodium Chloride Injection 0.9% (also may be used with Dextrose 5% and Sodium Chloride Injection 0.45% or 0.2%)
- Lactated Ringer's Injection
- Ringer's Injection
- Sodium Bicarbonate 5% in Sterile Water for Injection

Pharmacy Bulk Vial

Reconstitute according to the Pharmacy Bulk Vial Reconstitution Table. SHAKE TO DISSOLVE ALL POWDER. Further dilute aliquots in 50 to 100 mL of Sterile Water for Injection or one of the solutions listed above.

The further diluted solutions above should be used within 24 hours at room temperature (between 15 and 25°C) or 72 hours under refrigeration (between 2 and 8°C) from the time of initial puncture. Protect from light.

AVAILABILITY OF DOSAGE FORMS

Cefazolin for Injection USP (sterile cefazolin sodium) is supplied as a powder in 15 mL, glass vials, with rubber stoppers and flip-off caps, equivalent to 500 mg or 1 g of cefazolin, boxes of 10. Cefazolin for Injection USP is also available as a pharmacy bulk in a 100 mL infusion bottle, closed with hollow rubber stoppers and flip-off caps, equivalent to 10 g of cefazolin, box of 1.

THE AVAILABILITY OF THE PHARMACY BULK VIAL IS INTENDED FOR HOSPITALS WITH A RECOGNIZED IV ADMIXTURE PROGRAM.

LATEX-FREE STOPPER: Stopper contains no dry natural rubber.

MICROBIOLOGY

Cefazolin Activity Against Clinical Isolates

	No. of Strains		Cumulative Percentage Susceptible to Strains Indicated Concentration (mcg/mL)				
		<0.05	<0.1-0.78	1.56-3.13	6.25-12.5	25-50	100
<i>S. Aureus</i>	700	0.14	59.1	90.6 – 92.4*	97.3	99.7	99.9
<i>S. Pyogenes</i>	5	80+	100				
<i>S. Fæcalis</i>	2				50	100	
<i>S. Pneumoniæ</i>	6	100+					
<i>E. Coli</i>	484		8.7	67.9	92.1	95.9	97.7
<i>P. Mirabilis</i>	30			50	86.7	90	90
<i>K. Pneumoniæ</i>	138		2.9	53.6	73.2	91.3	93.5
<i>Enterobacter</i>	31			6.5	29.0	64.5	77.4
<i>H. Influenzæ</i>	30			13.3	70.0	100	
<i>N. Gonorrhoeæ</i>	13		38.5	100			
<i>Shigella SPP</i>	2			50	50	100	
<i>Salmonella SPP</i>	8			100			
<i>Staphylococci</i> (coagulase - negative)	295		66	82	90	93	100

* Reported as 3.13-6.25 mcg/mL

+ Reported as ≤0.1 mcg/mL

Disc Susceptibility Tests

The following criteria should be used to interpret tests using a standardized 30 mcg cephalosporin-class disc:

Zones of 18 mm or greater indicate that the tested organisms are susceptible and are likely to respond to therapy. Zones of 15 to 17 mm indicate organisms of intermediate susceptibility which may be susceptible if high dosage is used or if the infection is confined to tissues and fluids (e.g., urine) in which high antibiotic levels are attained. Zones of 14 mm or less are produced by resistant organisms.

The cephalothin disc should not be used for testing susceptibility to other cephalosporins.

Dilution Techniques: If the minimal inhibitory concentration (MIC) for cefazolin is not more than 16 mg/mL then a bacterial isolate may be considered susceptible. If the MIC is equal to or greater than 64 mg/mL, organisms are considered to be resistant.

The ranges of MIC for the control strains were:

E. Coli ATCC 25922 1.0-4.0 mg/mL

S. Aureus ATCC 25923 0.25-1.0 mg/mL

PHARMACOLOGY

Clinical Pharmacology

The blood levels of cefazolin listed on the following tables were determined following intramuscular and intravenous administration.

Serum Concentration (mg/mL) Following Administration

(Time After Intravenous Injection in Minutes)

	5	15	30	60	120	240
Cefazolin 1g	188.4	135.8	106.8	73.7	45.6	16.5

(Time After Intramuscular Injection in Hours)

Cefazolin	1/2	1	2	4	6	8
1g	65.8	68.3	60.6	29.3	11.2	6.5
500 mg	36.2	36.8	37.9	15.5	6.3	3.0
250 mg	15.5	17.0	13.0	5.1	2.5	<1.5

The serum half-life is approximately 1.8 hours following intravenous administration and 2.0 hours after intramuscular administration.

The mean peak serum levels of cefazolin in hospitalized patients are approximately equivalent to those seen in normal volunteers.

Healthy volunteers received a continuous intravenous infusion of 3.5 mg/kg for 1 hour (approximately 250 mg) and 1.5 mg/kg hourly for the next two hours (approximately 100 mg). A steady serum level of 28 mg/mL was attained at the third hour.

Cefazolin levels in synovial fluid and serum are similar four hours after drug administration. Levels in cord blood are equivalent to 40% of those found in maternal blood.

In patients without obstructive biliary disease, serum levels of cefazolin can be up to five times lower than bile levels of cefazolin. However, bile levels of cefazolin are considerably lower than serum levels in patients with obstructive biliary disease.

Cefazolin is excreted unchanged in the urine. Approximately 60% of the drug is excreted in the first six hours, and this increases to 70%-80% within 24 hours. Peak urine concentrations of approximately 2400 mcg/mL and 4000 mcg/mL are achieved following intramuscular doses of 500 mg and 1 gram, respectively.

TOXICOLOGY

Acute Toxicity

Parenteral and oral cefazolin demonstrated low toxicity in rodents, canines and rabbits tested in acute toxicity studies.

Acute Toxicity

Species	Route of Administration	LD ₅₀ (g/kg)
Mice	intravenous	≥3.9
	intraperitoneal	≥4.0
	subcutaneous	7.6
	oral	>11.0
Rats	intravenous	≥3.0
	intraperitoneal	7.4
	subcutaneous	>10
	oral	>11.0
Rabbits	intravenous	>2.0
Dogs	intravenous	>2.0

Subacute and Chronic Toxicity

Rats and dogs were studied in subacute and chronic parenteral toxicity of cefazolin. Rats were treated for 3 and 6 months subcutaneously and for one month intraperitoneally. The highest doses ranged from 2000 mg/kg per day in the 6 month study to 4000 mg/kg per day in the 1 and 3 month studies. Anemia was the only significant abnormality attributable to SC drug administration. In all experiments there was a definite dose-related depression of SGPT levels. Leukocytosis and hypererythropoiesis accompanied the anemia, which was probably related to hemorrhaging at the injection site.

The lowering of the SGPT was dependent upon both the dose and the duration of treatment. This was not statistically significant at the low doses and was reversible upon withdrawal of the drug. Equivalent chronic studies in dogs produced similar results: at the higher doses there was a fall in SGPT and frank anemia resulted from high subcutaneous doses. Dogs treated intravenously did not develop the anemia indicating that it was probably associated with hemorrhaging at the site of injection.

Reproduction and Teratology

Rabbits and mice were administered cefazolin in doses of 240 mg/kg/day and 2 400 mg/kg/day. No teratologic effects were observed. No adverse effects on mating, fertility, gestation, delivery and lactation were observed in rats administered 2000 mg/kg per day. Baby rats whose mothers were injected with 1200 mg/kg/day of cefazolin prior to delivery and throughout lactation were observed and there was no effect on the birth, or peri- and postnatal development.

Nephrotoxicity

The nephrotoxicity of cefazolin was studied following intravenous injections of rabbits and subcutaneous injections of mice and rats. The mean nephrotoxic intravenous dose in rabbits was between 300 and 400 mg/kg/day. No evidence of renal damage was produced when cefazolin was injected subcutaneously into mice at a dose of 8 g/kg/day for up to 3 days and into rats at a dose of 4 g/kg/day for up to 7 days.

REFERENCES

1. Birkhead HA, Briggs GB, Saunders LZ. Toxicology of cefazolin in animals. *J Infect Dis* 1973; 128:S379-S381.
2. Evrard J, Doyon F, Acar JF, et al. Two-day cefamandole *versus* five-day cephazolin prophylaxis in 965 total hip replacements. *Int Orthop* 1988; 12:69-73.
3. Handbook on Injectable Drugs, 7th Edition. Trissel LA (Ed). American Society of Hospital Pharmaceutics Inc. 1992.
4. Hemsell D, Hemsell P, Nobles B, et al. Moxalactam *versus* cefazolin prophylaxis for vaginal hysterectomy. *Am J Obstet Gynecol* 1983;147:379-385.
5. Ishiyarna S, Nakayama I, Iwamoto H, et al. Absorption, tissue concentration, and organ distribution of cefazolin. *Antimicrob Agents Chemother* 1970; 476-480.
6. Iversen P, Madsen PO. Complicated urinary tract infections treated with cefuroxime or cefazolin: A comparative study. *Clin Ther* 1981; 4(4):302-307.
7. Kaye D, Wenger N, Agarwal B. Pharmacology of intraperitoneal cefazolin in patients undergoing peritoneal dialysis. *Antimicrob Agents Chemother* 1978; 14(3):318-321.
8. Kini PM, Fernandez J, Causay RS, et al. Double-blind comparison of cefazolin and cephalothin in open-heart surgery. *J Thorac Cardiovasc Surg* 1978; 76:506-509.
9. Kirby WMM, Regamey C. Pharmacokinetics of cefazolin compared with four other cephalosporins. *J Infect Dis* 1973; 128:534-536.
10. Kozatani J, Okui M, Matsubara T, et al. Cefazolin, a new semisynthetic cephalosporin antibiotic VI. Excretion and metabolism of cefazolin-¹⁴C in rats after intramuscular administration. *J Antibiot* 1972; 25:86-93.
11. Lea AS, Sudan AW, Wood BA, et al. Randomized comparative study of moxalactam and cefazolin in the treatment of acute urinary tract infections in adults. *Antimicrob Ag Chemother* 1982; 22(1):32-35.
12. Levison ME, Levison SP, Ries K, Kaye D. Pharmacology of cefazolin in patients with normal and abnormal renal function. *J Infect Dis* 1973; 128-S354-S357.
13. Lorian V. Ed, *Antibiotics in Laboratory Medicine*, Second Edition, Williams & Wilkins, Baltimore, MD, pp. 1058-1059, 1986.
14. Mantel LA, Nicolle LE, Ronald AR, et al. A multicentre prospective randomized trial comparing ceftazidime with cefazolin/tobramycin in the treatment of hospitalized patients with non-pneumococcal pneumonia. *J Antimicrob Chemother* 1983; 12:S9-S20.

15. Marier RL, McCloskey RV, Dickenson G, et al. Comparative clinical trial of imipenem cilastatin (N-formimidoyl-thienamycindehydropeptidase inhibitor) and cefazolin. *J Antimicrob Chemother* 1983;12:133-139.
16. Mine Y, Nishida M. Cefazolin, a new semisynthetic cephalosporin antibiotic. I.V. Antigenicity of cefazolin and its cross reactivity with benzylpenicillin, ampicillin and cephaloridine. *J Antibiot* 1970; 23:195-203.
17. Nakano H. Cefazolin sodium, in Goldberg ME Ed, *Pharmacological and biochemical properties of drug substances*, American Pharmaceutical Association, Washington, DC, pp. 155-182, 1977.
18. Nishida M, Matsubara T, Murakawa T et al. Cefazolin, a new semisynthetic cephalosporin antibiotic. II. *In vitro* and *in vivo* antimicrobial activity. *J Antibiot* 1970; 23:137-148.
19. Nishida M, Matsubara T, Murakawa T, et al. Cefazolin, a new semisynthetic cephalosporin antibiotic. III. Absorption, excretion and tissue distribution in parenteral administration. *J Antibiot* 1970; 23:184-194.
20. Norrby SR. Side effects of cephalosporins. *Drugs* 1987; 34 (suppl.2):105-120.
21. Pankey GA, Katner HP, Valainis GT, et al. Overview of bacterial infections of the skin and soft tissue and clinical experience with ticarcillin plus clavulanate potassium in their treatment. *Am J Med* 1985; 79(suppl.2), 106-115.
22. Ram MD, Watanatittan S. Levels of cefazolin in human bile. *J Infect Dis* 1973; 128:S361-S363.
23. Regamey C, Gordon RC, Kirby WMM. Cefazolin vs cephalothin and cephaloridine. A comparison of their clinical pharmacology. *Arch Intern Med* 1974; 133: 407-410.
24. Reinartz JA, Kier CM, Guckian JC. Evaluation of cefazolin in the treatment of bacterial endocarditis and bacteremia. *J Infect Dis* 1973; 128:5392-5396.
25. Sabath LD, Wilcox C, Garner C, Finland M. *In vitro* activity of cefazolin against recent clinical bacterial isolates. *J. Infect Dis* 1973; 128: S320-S326.
26. Turck M, Clark RA, Beaty HN, et al. Cefazolin in the treatment of bacterial pneumonia. *J Infect Dis* 1973;138:5382-5385.
27. Wallace RJ, Martin RR, Quinones FJ. Ceforanide and cefazolin therapy of pneumonia: comparative clinical trial. *Antimicrob Ag Chemother* 1981;20:648-652.
28. Weinstein AJ. The cephalosporins: activity and clinical use. *Drugs* 1980;19:137-154.

29. Wick WE, Preston DA. Biological properties of three 3-heterocyclic-thiomethyl cephalosporin antibiotics. *Antimicrob Agents Chemother* 1972;1:221-234.
30. Teva Canada Limited, Cefazolin for Injection, Product Monograph, Control #212089, Date of Revision: July 30, 2018.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE PATIENT

MEDICATION INFORMATION

CEFAZOLIN FOR INJECTION USP Sterile cefazolin sodium powder

Read this carefully before you start taking Cefazolin for Injection USP and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Cefazolin for Injection USP.

What is Cefazolin for Injection USP used for?

Cefazolin for Injection USP is used for the treatment of infections caused by certain bacteria in many different parts of the body including the treatment of pneumonia.

Cefazolin for Injection USP can also be used to prevent infections, before and after surgery.

Antibacterial drugs like Cefazolin for Injection USP treat only bacterial infections. They do not treat viral infections.

How does Cefazolin for Injection USP work?

Cefazolin for Injection USP is an antibiotic, which belongs to a class of drugs called cephalosporins. Cefazolin for Injection USP works by killing bacteria which cause infections in the body.

What are the ingredients in Cefazolin for Injection USP?

Medicinal ingredients: cefazolin sodium sterile

Non-medicinal ingredients : none

LATEX-FREE STOPPER: Stopper contains no dry natural rubber.

Cefazolin for Injection USP comes in the following dosage forms:

Sterile powder for injection: 500 mg, 1 g and 10 g cefazolin per vial.

Do not use Cefazolin for Injection USP if:

- you have had an allergic reaction to Cefazolin for Injection USP or other medicines such as cephalosporins.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Cefazolin for Injection USP. Talk about any health conditions or problems you may have, including if you:

- have had an allergic reaction to penicillins
- have a history of bowel disease, particularly colitis
- have gallbladder problems
- have kidney problems with or without liver problems
- are pregnant or could become pregnant during treatment
- are breast feeding

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Cefazolin for Injection USP:

- Probenecid used in the treatment of gout

How to take Cefazolin for Injection USP:

- Cefazolin for Injection USP will be given to you by your healthcare professional as an injection into either a vein or a muscle.
- Although you may feel better early in treatment, Cefazolin for Injection USP should be used exactly as directed.
- Misuse or overuse of Cefazolin for Injection USP could lead to the growth of bacteria that will not be killed by Cefazolin for Injection USP (resistance). This means that Cefazolin for Injection USP may not work for you in the future.
- Do not share your medicine.

Usual dose:

Your healthcare professional will decide how much Cefazolin for Injection USP to give you and how often.

Overdose:

If you think you have been given too much CEFAZOLIN FOR INJECTION USP , contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss an appointment to receive an injection of Cefazolin for Injection USP, contact your healthcare professional as soon as possible.

What are possible side effects from using Cefazolin for Injection USP?

These are not all the possible side effects you may feel when taking Cefazolin for Injection USP. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- diarrhea, nausea, vomiting
- stomach cramps, loss of appetite
- rash, itching
- pain, tenderness or a hardened mass at the injection site
- vaginal and anal itching

Cefazolin for Injection USP can cause abnormal blood test results. Your healthcare professional will decide when to perform blood tests and interpret the results.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Anemia: fatigue, loss of energy, weakness, shortness of breath		√	
Hypersensitivity: rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			√
Liver disorder: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		√	
Oral candidiasis (yeast infection): creamy white bumps on the tongue, cheeks, gums or throat that bleed when scraped, pain, trouble swallowing, bad taste in the mouth		√	
Phlebitis: swelling of a vein near the injection site, with pain, tenderness, redness		√	
Platelet count increased: burning, redness, throbbing, numbness and/or tingling in the hands and feet, headache, dizziness, weakness, fainting, chest pain, vision changes		√	
Pseudomembranous colitis: watery, bloody diarrhea, mucus in the stool, abdominal cramps and pain, fever			√
Vulvovaginal mycotic infection: vaginal itching, burning during intercourse or urination, pain, redness, swelling, discharge		√	
White blood cell count decreased: infection, fatigue, fever, aches, pain, flu-like symptoms		√	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- d. Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- e. Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

How to store Cefazolin for Injection USP:

Cefazolin for Injection USP will be stored by your healthcare professional between 15°C and 25°C and protected from light.

Keep out of reach and sight of children.

If you want more information about Cefazolin for Injection USP:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website <https://www.canada.ca/en/health-canada.html>; the manufacturer's website <http://www.sandoz.com>; or by calling 1-800-361-3062

or by written request at:

110 rue de Lauzon
Boucherville QC
J4B 1E6

Or by e-mail at :

medinfo@sandoz.com

This leaflet was prepared by Sandoz Canada Inc.

Last revised: October 04, 2018.