

CONSUMER INFORMATION

Creo-Rectal®

Enfants/Children



WARNINGS

Keep out of reach of children. Read the complete instructions prior to use and follow them all. Do not use in children under 6 years of age. Overdose may result in serious harm. Do not exceed the single and maximum daily dose, because this will increase the risk of adverse drug reactions including convulsions and rectal burning. Do not use for longer than 3 days due to the possible risks related to the accumulation of eucalyptol and camphor in the body including the brain, in particular neuropsychological disorders; and the risk of rectal burning. Do not use in children with a history of epilepsy or febrile convulsions, or with a recent history of anorectal lesions. **DO NOT** give with any other cough and cold medications since harm may occur, unless recommended by a physician. Consult a physician prior to combining with other medications, including natural health products, prescription or nonprescription drugs. Consult a physician if symptoms worsen, last for more than a week or are accompanied by a high fever ($>38^{\circ}\text{C}$) or the production of thick yellow/green phlegm. Do not use if the child is allergic to any of the ingredients. Discontinue use if allergic reactions such as wheezing, rash or itching develop. Side effects may include: allergic reaction, breathing difficulties, convulsions, drowsiness, hallucinations, and rapid heart rate.

® Registered trademark owned/used under license by Sandoz Canada Inc.



Sandoz Canada Inc.
Qc, Canada J4B 7K8
medinfo@sandoz.com

☎1-800-361-3062

46112582